

Multi- k configurations

N. Bernhoeft,¹ J. A. Paixão,² C. Detlefs,³ S. B. Wilkins,^{4,3}

P. Javorsky,⁴ E. Blackburn,^{4,5} and G. H. Lander⁴

¹*département de Recherche Fondamentale sur la Matière Condensée,
CEA, F-38054 Grenoble CEDEX, France*

²*Physics Department, University of Coimbra, Coimbra, 3004-516 Portugal*

³*European Synchrotron Radiation Facility,
Boîte Postal 220, F-38043 Grenoble CEDEX, France*

⁴*European Commission, JRC, Institute for Transuranium Elements,
Postfach 2340, Karlsruhe, D-76125 Germany*

⁵*Institut Laue-Langevin, Boîte Postal 156, F-38042 Grenoble CEDEX, France*

(Dated: February 2, 2008)

Abstract

Using resonant x-ray scattering to perform diffraction experiments at the U M_4 edge novel reflections of the generic form $\langle kkk \rangle$ have been observed in $UAs_{0.8}Se_{0.2}$ where $\mathbf{k} = \langle k00 \rangle$, with $k = \frac{1}{2}$ reciprocal lattice units, is the wave vector of the primary (magnetic) order parameter. The $\langle kkk \rangle$ reflections, with 10^{-4} of the $\langle k00 \rangle$ magnetic intensities, cannot be explained on the basis of the primary order parameter within standard scattering theory. A full experimental characterisation of these reflections is presented including their energy, azimuthal and temperature dependencies. On this basis we establish that the reflections most likely arise from the electric dipole operator involving transitions between the core $3d$ and partially filled $5f$ states. The temperature dependence couples the $\langle kkk \rangle$ peak to the triple- $\mathbf{k}$ region of the phase diagram: Below ~ 50 K, where previous studies have suggested a transition to a double- $\mathbf{k}$ state, the intensity of the $\langle kkk \rangle$ is dramatically reduced. Whilst we are unable to give a definite explanation of how these novel reflections appear, this paper concludes with a discussion of possible ideas for these reflections in terms of the coherent superposition of the 3 primary (magnetic) order parameters.

PACS numbers: 75.25.+z, 75.10.-b

I. INTRODUCTION

Since their discovery in 1963 by Kouvel and Kasper¹, multi- $\mathbf{k}$ configurations have generated their share of confusion in the description of magnetic structures. The ambiguities arise since magnetic systems commonly lower their free energy by formation of domains. This eventuality frequently renders the best-known technique for their microscopic identification, neutron diffraction, impotent in the determination as to whether the magnetic structure is single- $\mathbf{k}$ or multi- $\mathbf{k}$. The respective characteristics are that a single- $\mathbf{k}$ configuration has only one magnetic propagation vector in any given magnetic domain whilst a multi- $\mathbf{k}$ configuration is defined by the *simultaneous* presence of more than one such propagation vector. Given the possibility of multiple scattering, one immediately sees the likelihood of confusion in the interpretation of diffraction peak intensities from a multi-domain sample.

Neutron diffraction is a bulk technique, sensitive to the spatial periodicities of the magnetic field modulation. In general one cannot locate the scattering volume from which the diffraction peaks arise to a precision better than that given by the incident and scattered beams' intersection with the sample. Given incident flux limitations, even at the most powerful neutron sources, beams can rarely be made sufficiently small (on the scale of magnetic domains) to identify unambiguously the magnetic configuration from intensity measurements in a multi-domain sample. External perturbations can, of course, change the domain populations and may allow identification, but this always begs the question as to whether the external perturbation may have changed the intrinsic magnetic configuration.

To be specific we take the case of the system $\text{UAs}_{1-x}\text{Se}_x$ where complete solid solutions exist and a considerable amount of neutron diffraction has been performed². Diagrams of the possible magnetic structures for $0 < x < 0.3$ are shown in Figure I. The single- $\mathbf{k}$ configuration has 3 distinct (tetragonal) domains, as does the $2\mathbf{k}$ state, whilst the cubic $3\mathbf{k}$ phase forms in a single domain. In these illustrations the repeat distance of the magnetic structure is twice the NaCl-type chemical unit cell, so the magnitude of the wave vector of the magnetic modulation is given by $k = \frac{1}{2}$ reciprocal lattice units (rlu). The primary magnetic reflections are then of the form $\langle k00 \rangle$ where the $\langle \dots \rangle$ indicates a permutation over indexes. These reflections, which are the *only* ones observed by neutron diffraction, are also imaged by resonant x-ray scattering (RXS), via the $F^{(1)}$ electric dipole (E1) scattering amplitude³. In addition, the RXS cross section exhibits $F^{(2)}$ dipole amplitudes^{4,5} which give




FIG. 1: $\text{UAs}_{1-x}\text{Se}_x$ crystallizes in the fcc NaCl structure. The magnetic modulation wavevector is $\langle k00 \rangle$ where $\langle \dots \rangle$ signifies all permutations of k , and in the composition discussed here $k = \frac{1}{2}$. The magnetic moments of the four uranium atoms within the chemical unit cell have the same direction and magnitude. For simplicity we have therefore shown only the magnetic moment in the corner of each chemical unit cell. (a) In a longitudinal single- $\mathbf{k}$ structure with $\mathbf{k} = [001/2]$, the moments are aligned along $[001]$ and the unit cell is doubled along this direction. (b) In the $2\mathbf{k}$ structure with $k_a = [100]$ and $k_b = [010]$, the net moment is along $[110]$, and the unit cell is doubled in the a and b directions. (c) In the $3\mathbf{k}$ structure the unit cell is doubled in all directions and the net moment direction is $[111]$. Of all the structures the $3\mathbf{k}$ is the only one in which a unique domain exists.

rise to peaks at positions of the form $\langle k k 0 \rangle$. The symmetries of the $F^{(1)}$ and $F^{(2)}$ terms may be exploited to distinguish respectively between single- $\mathbf{k}$ and $2\mathbf{k}$ or $3\mathbf{k}$ structures⁶.

Under the constraints of the geometric structure factor, the $F^{(2)}$ amplitude projects, on a given scattering centre, a *pair* of the $\langle k 0 0 \rangle$ order parameters which, given inter-site phase coherence yields Bragg diffraction peaks of the form $\langle k k 0 \rangle$. The respective $F^{(1)}$, $F^{(2)}$ assignments have been experimentally verified through the polarisation and azimuthal dependence of the scattered photons⁶. Moreover, even though both amplitudes are electric dipole in origin, the contributions from the $F^{(2)}$ uranium scattering amplitudes have different matrix elements and are distinguished by their resonant energy and lineshape from the $F^{(1)}$ profiles^{6,7,8}.

In the course of these experiments an additional group of reflections, much weaker than the other two sets described above, of the generic form $\langle k k k \rangle$ have been observed. In this paper we give details characterising these reflections and our difficulty in explaining them within conventional scattering theory.

II. EXPERIMENTAL DETAILS AND RESULTS

Experiments were performed with σ incident polarisation of the photon beam at the ID20 beamline⁹, ESRF, Grenoble, France in the configuration used in previous work⁶. The studies were carried out on a single crystal of $\text{UAs}_{0.8}\text{Se}_{0.2}$ which, above a tetragonal distortion at $T^* \sim 50$ K, exhibits the cubic rock salt structure³. On warming, $\text{UAs}_{0.8}\text{Se}_{0.2}$ is known to pass from a magnetic configuration of commensurate ($k = \frac{1}{2}$) to incommensurate ($k = 0.475$) wave vector at $T_0 \sim 119$ K, and to the paramagnetic state at $T_N \sim 124$ K^{2,3}. It has been shown, using a combination of both neutron and x-ray techniques^{2,6}, to adopt a multi- $\mathbf{k}$ structure for $T < 124$ K.

Representative reflections for the $\langle k 0 0 \rangle$, $\langle k k 0 \rangle$ and $\langle k k k \rangle$ peaks are illustrated in Figure 2, where the dependence on incident photon energy of the scattered intensity at the positions $(0\ 0\ 5/2)$, $(1/2\ 0\ 5/2)$ and $(1/2\ 1/2\ 5/2)$, for both the $\sigma \rightarrow \sigma$ and $\sigma \rightarrow \pi$ channels of the cross section at $T = 70$ K are shown. In all cases these energy scans are characterised by sharp wave vector profiles indicative of long-range order. As Figure 2 shows, the $(1/2\ 1/2\ 5/2)$ peak appears *only* in the $\sigma \rightarrow \pi$ channel with a resonant energy and width comparable to that of the $(0\ 0\ 5/2)$, evidence which already suggests that the $\langle k k k \rangle$ peak may arise from




FIG. 2: Reflections measured in a single crystal of $\text{UAs}_{0.8}\text{Se}_{0.2}$ with photons of energy near the $\text{U } M_4$ resonance, which is marked with a dashed vertical line. The temperature was 70 K for all panels and the wave vector is $k = \frac{1}{2}$. The incident polarization is σ , and using a $\text{Au}(111)$ analyser the open (closed) points correspond to intensity in the π (σ) channel. (a) The $\langle k00 \rangle$ reflection $(0\ 0\ 5/2)$ as observed in neutron diffraction. Note that it occurs only in the rotated $\sigma \rightarrow \pi$ channel (b) The reflection $(1/2\ 0\ 5/2)$ of the form $\langle kk0 \rangle$ is discussed in detail in Ref. 6. It arises because of the intrinsic non-collinearity of the $2\mathbf{k}$ magnetic structure. Note that in RXS contributions occur in both polarization channels, the energy maximum is shifted and the peaks are narrow (c) A new type of reflection $(1/2\ 1/2\ 5/2)$ of the form $\langle kkk \rangle$ as discussed in this paper. Intensity is only in the $\sigma \rightarrow \pi$ channel. The energy position and width are similar to case (a) above.




FIG. 3: Azimuthal scans in the $\sigma \rightarrow \pi$ channel about the scattering vector for the $(1/2 \ 1/2 \ 5/2)$ (open points) and $(-1/2 \ 1/2 \ 5/2)$ (closed points) reflections. The lines correspond to the analysis described in the text. Note that for all azimuthal angles the intensity in the scattered $\sigma \rightarrow \sigma$ channel is zero.

the $F^{(1)}$ dipole (E1) amplitude.

Figure 3 shows the azimuthal dependence of the intensity of the $\langle kkk \rangle$ reflections $(1/2 \ 1/2 \ 5/2)$ and $(-1/2 \ 1/2 \ 5/2)$ in the $\sigma \rightarrow \pi$ channel. The smooth variation of the intensity eliminates multiple scattering as a possible source of these peaks. The lines are calculated from the $F^{(1)}$ term of the E1 cross section, assuming a symmetry breaking vector along $\langle 111 \rangle$, parallel to $\langle kkk \rangle$ {i.e. along $[1 \ 1 \ 1]$ for the $(1/2 \ 1/2 \ 5/2)$ and along $[-1 \ 1 \ 1]$ for the $(-1/2 \ 1/2 \ 5/2)$ }. The agreement between the data and this model with only *one* overall




FIG. 4: Peak intensity of the $\langle k00 \rangle$, $\langle kk0 \rangle$ and $\langle kkk \rangle$ (circles, squares and diamonds respectively) as a function of temperature.

scale factor is excellent. These aspects are discussed further below.

The temperature dependencies of the $(0\ 0\ 5/2)$, $(1/2\ 0\ 5/2)$ and $(1/2\ 1/2\ 5/2)$ peaks are given in the Figure 4. The $(0\ 0\ 5/2)$ reflection represents one primary order parameter whilst the $2\mathbf{k}$ $(1/2\ 0\ 5/2)$ reflection involves two simultaneously present at each scattering centre and propagating with fixed phase relationship. The temperature dependencies of $I_{\langle k00 \rangle}$, $I_{\langle kk0 \rangle}$ and $I_{\langle kkk \rangle}$ for $70\text{ K} < T < 117\text{ K}$ in Figure 2 lie approximately in the ratio $T^{0.4} : T : T^{1.5}$.

Detailed specific heat and magnetisation measurements were made on the sample used in the diffraction experiments at the User Facility, Institute for Transuranium Elements, Karlsruhe¹⁰. We show in Figure 5 the results of (a) the integrated intensity of the $(1/2\ 1/2\ 5/2)$ peak, (b) heat capacity at zero field, and (c) the susceptibility in a magnetic field of 0.1 T, as a function of temperature. The loss of intensity observed in the $\langle kkk \rangle$ reflection is coupled to the nominal $2\mathbf{k} \Leftrightarrow 3\mathbf{k}$ transitions seen in both specific heat and susceptibility at T^* (Ref. 3).




FIG. 5: Integrated intensity of the $(\frac{1}{2}\frac{1}{2}\frac{5}{2})$ reflection in the region $35 \text{ K} < T < 60 \text{ K}$ (top panel), heat capacity at zero applied magnetic field (middle panel) and magnetic susceptibility in a applied magnetic field of 0.1 T (bottom panel).

III. DISCUSSION

Before turning to a possible explanation of this effect, we recapitulate the phase diagram of the $\text{UAs}_{0.8}\text{Se}_{0.2}$ as suggested by earlier neutron² and RXS experiments³. First, the disap-

pearing intensity of the $\langle kkk \rangle$ peak and the presence of a lattice distortion below $T^* \sim 50$ K³, show that the low-temperature state is most likely a $2\mathbf{k}$ phase². Above T^* , high-resolution x-ray experiments have not been able to detect any distortion from cubic symmetry, this suggests, in agreement with the neutron results, that the sample is not in a simple $2\mathbf{k}$ (or single- $\mathbf{k}$) phase. Previous authors have suggested this is be a $3\mathbf{k}$ state².

We now examine possible origins of the $\langle kkk \rangle$ peaks. A simple explanation would be that at the level of 10^{-4} of the total volume there are regions that exhibit an ordering at a single- $\mathbf{k}$ wave vector $\langle \frac{1}{2} \frac{1}{2} \frac{1}{2} \rangle$. This would explain the observed energy and azimuthal dependence of the scattering, Figures 2 and 3, respectively. However, there are at least two observations which contradict such a scenario. First, the similar, sharp, $\mathbf{q}$ widths of $\langle k00 \rangle$, $\langle kk0 \rangle$ and $\langle kkk \rangle$ reflections are indicative that the $\langle kkk \rangle$ peaks represent (bulk) long-range order. Second, the simple relation of their temperature dependencies to the $\langle k00 \rangle$ and $\langle kk0 \rangle$ peaks for $T^* < T < T_o$ would then have to be completely fortuitous, which is hard to accept. Moreover, single- $\mathbf{k}$ $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ ordering has, to date, never been reported in NaCl-structure uranium compounds. These observations all suggest that the $\langle kkk \rangle$ reflections are intimately related to the primary long-range order parameters of the material.

As already noted, both the electric dipole crosssection and the geometric structure factor of the magnetic moment (axial vector) for $\mathbf{Q} = \langle kkk \rangle$ vanish. The lowest combination of order parameters with finite geometrical structure factors is of rank 3, i.e. of the form $J_x J_y J_z$. For example, a symmetrised, octupolar operator couples directly to the $F^{(3)}$ term of the E2 cross section⁵, as observed e.g. in V_2O_3 ^{11,12}. To date however, there is no evidence of any E2 resonances at the actinide $M_{4,5}$ edges, since these would couple to the g states, with a correspondingly small matrix element. Furthermore, the E2 cross section would give rise to scattering in both the $\sigma \rightarrow \sigma$ and $\sigma \rightarrow \pi$ polarization channels, and one would expect the maximum of the resonance to be shifted towards the pre-edge region, as observed in transition metal¹¹ and rare earth systems^{13,14}. Rather, the energy and polarization profiles link these reflections to the $F^{(1)}$ term of the E1 cross section.

The effective moment direction along $\langle 111 \rangle$, as indicated by analysis of the azimuthal dependencies shown in Figure 3, indicates the origin of this resonance lies in the atomic matrix elements. Any combined lattice distortion or charge density wave (CDW) at $\langle kk0 \rangle$ with a magnetic dipole $\langle k00 \rangle$ construction is not supported by our observations. Furthermore, there is no experimental evidence for either a distortion or CDW in the cubic phase

for such hypothetical constructions¹⁵.

A mechanism to couple an E1 resonance to the octupolar moment of the valence shell was recently suggested by Lovesey et al.⁸. They found that the rank-2 tensor (E1-F⁽²⁾ term in the cross section) observed in NpO₂ may exist even in the absence of a quadrupolar moment on the Np ion. This tensor is constructed from a magnetic octupole (rank 3) and an induced Zeeman splitting in the 3d core shell (rank 1). The product of these tensors contains one of rank 2 which may have been observed in the x-ray experiment⁷.

In a similar framework coupling a rank (R) and rank ($R + 1$) tensor will yield a vector (rank 1) in its product yielding a $\langle kkk \rangle$ scattering amplitude. Hypothetical examples include combining a $\langle k00 \rangle$ magnetic moment in the 5f valence level with a $\langle kk0 \rangle$ quadrupolar splitting of the 3d core levels or vice versa. A still more complex scenario would be a $J_x J_y J_z$ octupolar moment in the 5f shell (rank 3) with a rank 2 quadrupolar splitting of the 3d core levels. What physical field would give the required core level anisotropy is unknown and such high level mechanisms are currently without foundations.

We conclude with a brief discussion of the phase diagram. Due to the absence of any measurable tetragonal distortion it was previously suggested that the phase between T^* and T_O is a $3\mathbf{k}$ state. However, there is no direct evidence for this assumption. In fact unpublished field-dependent specific heat measurements, as well as other results are rather difficult to interpret on this basis. Thus, the nature of the transition at T^* remains unclear. We suggest the low-temperature (tetragonal) configuration to be a state of $2\mathbf{k}$ domains composed of phase coherent pairs of the primary order parameters, in agreement with earlier work^{2,3}. However, as the temperature is raised, the tetragonal $2\mathbf{k}$ phase melts with fluctuations of increasing frequency between the possible domains giving, above T^* , a dynamic state which maintains a cubic environment within which both $2\mathbf{k}$ and $3\mathbf{k}$ correlations of the primary order parameters may coexist. At present we have no direct evidence on the lifetime of either $2\mathbf{k}$ or $3\mathbf{k}$ correlations above T^* . The rapid nature of the RXS technique (temporal resolution of $\sim 10^{-15}$ s) may be of importance if the coherence time scales are on the scale of the inverse bandwidth.

Whilst this paper eliminates many of the most obvious explanations for the presence of the $\langle kkk \rangle$ reflections clearly more work is needed to explain their observation at both *qualitative* and *quantitative* levels. We hope the observations and discussion will stimulate further experiments and theoretical studies of such multi- $\mathbf{k}$ systems.

Acknowledgments

We thank Ted Forgan, Christian Vettier, Anne Stunault, and François de Bergevin for interesting discussions and Matt Longfield for his help in the early part of these investigations. SBW, PJ and EB would like to thank the European Commission for support in the frame of the ‘Training and Mobility of Researchers’ programme.

-
- ¹ J. S. Kouvel and J. Kasper, *J. Phys. Chem. Solids* **24**, 529 (1963).
- ² M. Kuznietz, P. Burlet, J. Rossat-Mignod, and O. Vogt, *J. Magn. Magn. Matt.* **69**, 12 (1987).
- ³ M. J. Longfield, W. G. Stirling, E. Lidström, D. Mannix, G. H. Lander, A. Stunault, G. J. McIntyre, K. Mattenberger, and O. Vogt, *Phys. Rev. B* **63**, 134401 (2001).
- ⁴ J. P. Hannon, G. T. Trammell, M. Blume, and D. Gibbs, *Phys. Rev. Lett.* **61**, 1245 (1988).
- ⁵ J. P. Hill and D. F. McMorrow, *Acta Crystallogr. A* **52**, 236 (1996).
- ⁶ M. J. Longfield, J. A. Paixão, N. Bernhoeft, and G. H. Lander, *Phys. Rev. B* **66**, 054417 (2002).
- ⁷ J. A. Paixão, C. Detlefs, M. J. Longfield, R. Caciuffo, P. Santini, N. Bernhoeft, J. Rebizant, and G. H. Lander, *Phys. Rev. Lett.* **89**, 187202 (2002).
- ⁸ S. W. Lovesey, E. Balcar, C. Detlefs, G. van der Laan, D. S. Silvia, and U. Staub, *J. Phys.: Condens. Matt.* **15**, 4511 (2003).
- ⁹ <http://www.esrf.fr/UsersAndScience/Experiments/XASMS/ID20/>.
- ¹⁰ <http://itu.jrc.cec.eu.int/>.
- ¹¹ L. Paolasini, C. Vettier, F. de Bergevin, F. Yakhov, D. Mannix, A. Stunault, W. Neubeck, M. Altarelli, M. Fabrizio, P. A. Metcalf, et al., *Phys. Rev. Lett.* **82**, 4719 (1999).
- ¹² S. W. Lovesey and K. S. Knight, *J. Phys.: Condens. Matt.* **12**, L367 (2000).
- ¹³ D. Gibbs, G. Grübel, D. R. Harshman, E. D. Isaacs, D. B. McWhan, D. Mills, and C. Vettier, *Phys. Rev. B* **43**, 5663 (1991).
- ¹⁴ K. Hirota, N. Oumi, T. Matsumura, H. Nakao, Y. Wakabayashi, Y. Murakami, and Y. Endoh, *Phys. Rev. Lett.* **84**, 2706 (2000).
- ¹⁵ V. E. Dmitrienko and E. N. Ovchinnikova, *Acta Crystallogr. A* **56**, 340 (2000).